(C) at least one opioid antagonist; said dosage form further comprising a sustained

release carrier which causes said opioid agonist to be released over a time period of about 8 to about 24 hours when orally administered to a human patient.

3. (Amended) The oral dosage form of claim 1, wherein the opioid agonist is hydrocodone or a pharmaceutically acceptable salt thereof and the antagonist is naltrexone or a pharmaceutically acceptable salt thereof.

- 9. (Amended) The oral dosage form of claim 6, wherein said opioid antagonist is selected from the group consisting of naltrexone, naloxone, nalmephene, cyclazocine, levallorphan, pharmaceutically acceptable salts thereof and mixtures thereof.
- 10. (Amended) The oral dosage form of claim 6, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof.
- 12. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is oxycodone or a pharmaceutically acceptable salt thereof.
- 13. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is codeine or a pharmaceutically acceptable salt thereof.
- 14. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is hydromorphone or a pharmaceutically acceptable salt thereof.

- 15. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is levorphanol or a pharmaceutically acceptable salt thereof.
- 16. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is meperidine or a pharmaceutically acceptable salt thereof.
- 17. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is methadone or a pharmaceutically acceptable salt thereof.
- 18. (Twice Amended) The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone or a pharmaceutically acceptable salt thereof and said opioid agonist is morphine or a pharmaceutically acceptable salt thereof.
- 29. (Amended) The oral dosage form of claim 28 wherein said opioid antagonist is selected from the group consisting of naltrexone, naloxone, nalmephene, cyclazocine, levallorphan, pharmaceutically acceptable salts thereof and mixtures thereof.
- 32. (Thrice Amended) A method of treating pain, comprising administering an oral dosage form according to claim 1 orally to a human patient in an analgesically effective amount.